

SEQUENCE LISTING

<110> Rothschild, Max F. Vincent, Amy L. Tuggle, Christopher K. <120> PROLACTIN RECEPTOR GENE AS A GENETIC MARKER FOR INCREASED LITTER SIZE IN ANIMALS <130> P02285UD5 <140> US 09/900,063 <141> 2001-07-06 <150> US 08/812,208 <151> 1997-03-06 <150> US 08/742,805 <151> 1996-11-01 <150> US 60/022,180 <151> 1996-07-19 <160> 18 <170> PatentIn version 3.2 <210> 1 <211> 20 <212> DNA <213> Porcine 20 <400> 1 cccaaaacag caggagaacg <210> 2 <211> 20 <212> DNA <213> Porcine 20 <400> 2 ggcaagtggt tgaaaatgga <210> 3 <211> 492 <212> DNA <213> Porcine <220> <221> allele <222> (240)..(240) <223> r = a or g <400> 3

```
aagtcaacaa agatggagca ctggcgttgc tcccaaaaca gcaggagaac ggcgaccggc
                                                                    60
cggagaaggc tggcgccct gaaaccagca aggaatacgc ccaggtgtcc cgggtgatgg
                                                                    120
ataaccacat cctggtgtta gtgcaggatc cgcgagctcg aaacgtggct ccgtttgaag
                                                                    180
aaccaaccaa ggagaccccg ccatcccggc cgcagaatcc agctgcgaaa gacctggccg
                                                                    240
gcttcaccac ggccccgggc cactgcagac acccgctggg tgggctggat tacctcgatc
                                                                    300
ccgcaggctt tatgcactcc tttcagtgag agcttggttc atgggatgat gggttacaag
                                                                     360
gtggggtttt tttcaggtcg cactacgtga aatgcactct accagagaaa gctcgaaaat
                                                                     420
ggggttagaa tgacactacc cagactcaca gttcactcct cttcatgctc cattttcaac
                                                                     480
                                                                     492
cacttgcctc tt
<210> 4
 <211> 20
 <212> DNA
 <213> Unknown
 <220>
 <223> Human and rabbit cDNA sequences encoding the prolactin receptor
        were used to design the degenerate primer
 <220>
 <221> misc_difference
 <222> (13)..(13)
 <223> y = c or t
 <400> 4
                                                                       20
 tcacaaggtc aacaaagatg
 <210> 5
 <211> 20
 <212> DNA
 <213> Unknown
  <223> Human and rabbit cDNA sequences encoding the prolactin receptor
        were used to design the degenerate primer
```

<220>

<221> misc_feature

<223> r = g or a

<220>

<221> misc_difference

<222> (3)..(3)

<223> r = g or a

<220> <221> misc_difference <222> (9)(9) <223> r = g or a	
<220> <221> misc_difference <222> (16)(16) <223> r = g or a	
<400> 5 tggagaaaga ggcaagtggt	20
<210> 6 <211> 20 <212> DNA <213> Porcine	
<400> 6 cggccgcaga atcctgctgc	20
<210> 7 <211> 24 <212> DNA <213> Porcine	
<400> 7 accccacctt gtaacccatc atcc	24
<210> 8 <211> 18 <212> DNA <213> Porcine	
<400> 8 caaggtggga acatgagt	18
<210> 9 <211> 20 <212> DNA <213> Porcine	
<400> 9 aatcctttta tttttggccc	20
<210> 10 <211> 20 <212> DNA <213> Porcine	
<400> 10 gattattgtc tgggcagtgg	20

<210> 11 <211> 20 <212> DNA <213> Porcine	
<400> 11 aatcctttta tttttggccc	20
<210> 12 <211> 20 <212> DNA <213> Porcine	
<400> 12 gattattgtc tgggcagtgg	20
<210> 13 <211> 23 <212> DNA <213> Porcine	
<400> 13 ctatttcaca aaaactgcgc tac	23
<210> 14 <211> 566 <212> DNA <213> Porcine	
<220> <221> misc_feature <222> (36)(36) <223> n = unknown nucleotide position 36	
$<\!400\!>-14$ gattattgtc tgggcagtgg tettetetgt etatenacee eecteecatt catggetete	60
agggtataat ggccaaaaaa aagacaagac aaaaatgatg gaaacctaca gataattyaa	120
gcacctcatt ttgccattag ctgcattagc cataaaaaaa aaaaaaaaaa	180
gtgctagaaa aaaacagaat agactcattt gaaactgatc ttctctctac caaagggagt	240
agcgcagttg tgaaatagta aacgtctgac aagaacagca aataatccca ctagtaattt	300
cagaatccgc ctcctcaatt agccagaatt cactgtgatg ctggcctcta taattattat	360
ttgtcttcac cactgattag tttcacatca tgaaaattgc atgtcattta gtttcacrta	420
gesteagaas caaccetaat tectacetge catatecetg tageagetat tegaagatea	480

caaggtggga	acatgtgtya	tttatctttt	ctcttacatt	attttagagc	atggtggcct	540
gcatccgggc	caaaaataaa	aggatt				566
<210> 15 <211> 518 <212> DNA <213> Por						
<222> (49	c_feature)(49) unknown in	nucleotide	position 4	9		
<222> (67	c_feature)(67) unknown in	nucleotide	position 6	7		
<400> 15	acacacacac	acacacacac	acacaccacc	gttaagctnt	ctttctgaat	60
					gcacttctga	120
					ccccacccat	180
					tgactccagt	240
					caaatgtcgt	300
tctcccgaa	a aggaaacatt	cgcctgctg	g tggaagccgg	g gggcggatgq	g aggacttcct	360
accaactag	a cgctgactta	a ccacaaggaa	a gggtaagcat	t tegegtgtet	t cccaacaaac	420
					a gaactgcctc	480
	t tecegteat					518
<210> 16 <211> 13 <212> DN <213> Ho	3					
<400> 16	ic tectagaaa	a cctgagatc	t ttaaatgtc	g ttctcccaa	t aaggaaacat	60
					t tcactgactt	120
accacaggg		- 550				133
accacayys	,∽ ∽33					
<210> 17 <211> 53						

<212> PRT

<213>	Н	omo	sapi	ens												
<400>																
Gly Gl	ln	Leu	Pro	Pro 5	Gly	Lys	Pro	Glu	Ile 10	Phe	Lys	Ser	Pro	Phe 15	Cys	
Arg Se	er	Pro	Asn 20	Lys	Glu	Thr	Phe	Thr 25	Glu	Ala	Arg	Pro	Gly 30	Thr	Asp	
Gly G	ly	Leu 35	Pro	Thr	Asn	Lys	Pro 40	Ala	Asn	Tyr	Ser	Leu 45	Thr	Tyr	His	
Arg G 5	lu 0	Gly	Thr	Lys												
<210>		18														
<211><212><213>	•		ine													
<400>	> ca	18 cct	ccac	ggaaa	ac c	tgag	gatct	t ca	aatg	gtcgt	tct:	cccg	jaaa	agga	aacatt	60
															gactta	
ccaca																132